

Features:

- Enzyme with hot start capability increases reaction specificity and sensitivity, optimized
- DFS-Taq PLUS DNA polymerase activation requires not more than 5 min heating
- High selectivity and reaction yield
- The mix is colored for easy pipetting
- Reduced preparation time

Applications:

- Real-time PCR with intercalating dye SYBR Green I and ROX as reference dye
- Conventional PCR
- High-throughput PCR
- Genotyping
- For detection of bacterial DNA

Platforms:

(ABI) 7500, 7500 Fast, ViiA 7, QuantStudio 12K; Stratagene Mx4000, Mx3005P, and Mx3000P

Description:

Bio-Star qPCR-Mastermix SYBRGreen LOW-ROX (2x) is developed for quantitative real-time PCR with fluorescent dye SYBR Green I. The Mastermix contains all components, except template and primers, for successful PCR.

- The mix is optimized for efficient and reproducible hot-start real-time PCR of genomic, plasmid and viral DNA samples.
- The solution contains substances that increase half-life and processivity of DFS-Taq PLUS DNA polymerase by enhancing its stability during PCR.
- It includes components that influence primer annealing temperature and characteristics of template melting thus enabling to increase the specificity of PCR and use templates with complicated structure.
- DFS-Taq Plus DNA polymerase is inactive at room temperature because of monoclonal antibodies.
- The inert dye allows control when using multi-well plates. Use of the kit saves time and minimizes contamination risk due to reduced number of pipetting steps.

Components and Mixture

2X Bio-Star qPCR-Mastermix SYBRGreen LOW-ROX contains:

- 100 mM Tris-HCl (pH 8.5 at 25 °C), 100 mM KCl, 0.4 mM each of dNTP, 3 mM MgCl₂, 0.06 U/μl DFS-Taq DNA Polymerase, 0.025% Tween 20, stabilizers and enhancers, SYBR Green I, 60 nM ROX fluorescent dye, and inert dye
- Tube of MgCl₂ 100 mM
- Water Mol.Bio Grade 2 x 1,25 ml

Storage and transportation: at -20 °C; not more than 50 thawing-freezing cycles. shipping with blue ice or at room temperature

Storage terms: up to 24 months

Stability tests / Quality control / Comparison

Exodeoxyribonuclease activity: DNA was stable after incubation of 1 µg fragment of phage lambda DNA in the presence of 25 µl of *Bio-Star qPCR Mastermix (2x)* in 50 µl reaction solution at 37 °C and 70 °C for 4 h.

Test for bacterial DNA contamination

Test for Hot -Start (we test activity with or without MAB at various temperatures

Exodeoxyribonuclease activity: DNA was stable after incubation of 1 µg fragment of phage lambda DNA in the presence of 25 µl of BioMaster HS-qPCR (2x) in 50 µl reaction solution at 37 °C and 70 °C for 4 h.

Ribonuclease activity: Absence of ribonuclease activity was confirmed after incubation of 1 µg of 5'-[P32]-labeled RNA fragment in the presence of 25 µl of BioMaster HS-qPCR (2x) in 50 µl reaction solution at 37 °C for 4 h.

Stability: One month at room temperature does not reduce PCR efficiency

Amplification protocol

1. Defrost the reaction mixture and stir thoroughly.
2. Add the following components into the thin-wall PCR tubes considering the final volume of a reaction mixture equal to 50 µl:

Component	Volume	Final concentration
2x Mastermix	25 µl	1x
Forward Primer	variable	0.1 - 600 nM
Reverse Primer	variable	0.1 - 600 nM
DNA Template	variable	10 pg - 1 µg
Sterile Water	up to 50 µl	

3. Gently vortex and remove droplets by centrifugation.
4. Perform PCR using temperature conditions recommended below:

Step	Temp.°C	Incubation time	Number of Cycles
Preliminary denaturation	95	3-7 min	1
Denaturation	95	15 sec	25-40
Annealing	50-68	10-30 sec	25-40
Elongation	58-72	30-60 sec	25-40
Melting curve (recommended)	65-95		1

Order Information

Prod. No.	Description	Quantity
Y9250	Bio-Star qPCR-Mastermix SYBRGreen LOW-ROX (2x)	100 rcs (2,5 ml)